

Refine Search

Search Results -

Terms	Documents
triethanolamine adj10 (oleanolic or ursolic or betulinic)	3

Database:

US Pre-Grant Publication Full-Text Database
US Patents Full-Text Database
US OCR Full-Text Database
EPO Abstracts Database
JPO Abstracts Database
Derwent World Patents Index
IBM Technical Disclosure Bulletins

Search:

L2



Refine Search

Recall Text



Clear

Interrupt

Search History

DATE: Monday, October 22, 2007 [Purge Queries](#) [Printable Copy](#) [Create Case](#)

Set Name Query

side by side

Hit Count Set Name

result set

*DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; PLUR=YES; OP=OR*L2 triethanolamine adj10 (oleanolic or ursolic or betulinic) 3 L2L1 triethanolamine adj5 (oleanolic or ursolic or betulinic) 3 L1

END OF SEARCH HISTORY

[First Hit](#)[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

Generate Collection

Print

L3: Entry 4 of 15

File: PGPB

Aug 22, 2002

DOCUMENT-IDENTIFIER: US 20020115609 A1

TITLE: Materials and methods for making improved micelle compositions

Detail Description Paragraph:

[0128] Results indicated that progesterone (essentially insoluble in water as discussed above) in 10 mg/ml DSPE-PEG was soluble up to 198.5 .mu.g/ml. This result was consistent with results using betulinic acid, sparingly soluble in water (see Merck Index, 12th Edition, p. 1213), which was soluble up to 200 .mu.g/ml in 10 mg/ml DSPE-PEG. In similar experiments with betulinic acid (also insoluble as defined in USP), solubility was calculated at 250 .mu.g/ml in either SSM or SSC.

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

[First Hit](#) [Fwd Refs](#)[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

Generate Collection

Print

L3: Entry 7 of 15

File: USPT

Jan 3, 2006

DOCUMENT-IDENTIFIER: US 6982284 B1

TITLE: Compositions and methods for modification of skin lipid content

Brief Summary Text (50):

As discussed below, because ursolic acid compounds are highly insoluble in many solvents, including water, administration of such compounds in liposomes is a particularly preferred embodiment of the invention.

Description Paragraph (17):

Liposomal formulations are preferred because ursolic acid is highly insoluble in many solvents, particularly water, and common emulsifiers such as LECINOL S-10 have little effect. In accordance with the invention, this insolubility problem is addressed by taking advantage of the flat, planar structure of ursolic acid to stack it between the lipid tails in the phospholipid bilayer membranes of liposomes. Due to the charged headgroup of the phospholipids, liposomes containing ursolic acid are readily soluble in water.

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

Refine Search

Your wildcard search against 10000 terms has yielded the results below.

Your result set for the last L# is incomplete.

The probable cause is use of unlimited truncation. Revise your search strategy to use limited truncation.

Search Results -

Terms	Documents
(ursolic or betulinic or oleanolic) adj15 (alkal\$ adj3 solub\$)	1

Database:

US Pre-Grant Publication Full-Text Database
 US Patents Full-Text Database
 US OCR Full-Text Database
 EPO Abstracts Database
 JPO Abstracts Database
 Derwent World Patents Index
 IBM Technical Disclosure Bulletins

Search:

L6

Refine Search

Recall Text

Clear

Interrupt

Search History

DATE: Monday, October 22, 2007 [Purge Queries](#) [Printable Copy](#) [Create Case](#)

Set Name Query

side by side

Hit Count Set Name

result set

DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; PLUR=YES; OP=OR

<u>L6</u>	(ursolic or betulinic or oleanolic) adj15 (alkal\$ adj3 solub\$)	1	<u>L6</u>
<u>L5</u>	(ursolic or betulinic or oleanolic) adj10 (alkal\$ adj3 solub\$)	1	<u>L5</u>
<u>L4</u>	(ursolic or betulinic or oleanolic) adj5 (alkal\$)	17	<u>L4</u>
<u>L3</u>	(ursolic or betulinic or oleanolic) adj5 (insoluble or sol\$)	15	<u>L3</u>
<u>L2</u>	(ursolic or betulinic or oleanolic) same liposome same base	0	<u>L2</u>
<u>L1</u>	(ursolic or betulinic or oleanolic) same liposome	116	<u>L1</u>

END OF SEARCH HISTORY